



Working to Change the Trajectory of Lung Cancer

At Johnson & Johnson, we are focused on tackling lung cancer from every angle and are driven by our passion to help patients.

The Impact of Lung Cancer¹

Highest global death rate of all cancers

2nd most common cancer globally, behind only breast cancer

On average, 5,000 lives are claimed every day

Lung cancer is difficult to diagnose early.

- ↓ It starts growing very **deep in the lung**²
- ↓ Symptoms only start when it has grown significantly and **spread outside the lung**²

Early diagnosis is key.

- ↓ Earlier diagnosis could mean easier treatment and potentially cure³
- ↓ But **most people are diagnosed at advanced stages**, when treatment options are limited³

Screening is critical for those at increased risk.

- ↓ Regular **low-dose CT (LDCT) scans are the most important step** at-risk people can take toward early detection²
- ↓ Risk factors can include smoking, genetic susceptibility, poor diet, occupational exposures, and air pollution⁴

Anyone can get lung cancer.

- ↓ Up to **25% of lung cancers worldwide occur in people who have never smoked**⁵
- ↓ Stigma or blame are common, preventing timely screening and treatment⁶



A multifaceted disease requires a multidisciplinary approach.

Innovation at Johnson & Johnson has helped pave the way in oncology for more than 30 years. We are working holistically across the company to drive early detection, treatment and ongoing research and development **to help make a difference in the lives of millions of people.**

Maintaining health and creating a sense of hope for all people impacted by lung cancer are what drive us. **Together, we aim to change the trajectory of lung cancer.**



REFERENCES

¹ National Institutes of Health. CT screening significantly reduces lung cancer mortality. Available at: <https://www.nih.gov/news-events/nih-research-matters/ct-screening-significantly-reduces-lung-cancer-mortality>. Last accessed: July 2022.

² International Agency for Research on Cancer. GLOBOCAN 2020: Lung. Available at: <https://gco.iarc.fr/today/data/factsheets/cancers/15-Lung-fact-sheet.pdf>. Last accessed: July 2022.

³ Couraud S, Zalcman G, Milleron B, et al. Lung cancer in never smokers – A review. Eur J Cancer. 2012;48(12): 1299-1311. <https://www.sciencedirect.com/science/article/abs/pii/S0959804912002523>. Last accessed: July 2022.

⁴ Scott N, Crane M, Lafontaine M, et al. Stigma as a barrier to diagnosis of lung cancer: patient and general practitioner perspectives. Prim Health Care Res Dev. 2015;16(6):618-622. <https://www.cambridge.org/core/journals/primary-health-care-research-and-development/article/stigma-as-a-barrier-to-diagnosis-of-lung-cancer-patient-and-general-practitioner-perspectives/00DCD12CBBDCBC4517DCD678924BC277>. Last accessed: July 2022.

⁵ National Cancer Institute. SEER cancer stat facts: Lung and bronchus cancer. Available at: <https://seer.cancer.gov/statfacts/html/lungb.html>. Last accessed: July 2022.

⁶ Malhotra J, Malvezzi M, Negri E, et al. Risk factors for lung cancer worldwide. Eur Respir J. 2016;48(3):889-902. <https://erj.ersjournals.com/content/48/3/889>. Last accessed: July 2022.